



## Information about the Screening Checklist for Contraindications to Inactivated Injectable Influenza Vaccination (IIV or RIV)

### 1. Did you receive the flu vaccine last year?

This question helps us keep our records up to date and assists us in determining the number of vaccines a child may need during the current flu season.

If the patient indicates the flu vaccine was administered at another facility please verify it is documented in the chart.

### 2. Has the person to be vaccinated ever had a serious reaction to influenza vaccine in the past?

Patients reporting a serious reaction to a previous dose of inactivated influenza vaccine should be asked to describe their symptoms. Immediate – presumably allergic – reactions are usually a contraindication to further vaccination against influenza.

Fever, malaise, myalgia, and other systemic symptoms most often affect people who are first-time vaccines. These mild-to-moderate local reactions are not a contraindication to future vaccination. These people can receive injectable vaccine without further evaluation.

### 3. Do you have an allergy to a component of the vaccine (including eggs)?

All vaccines, including influenza vaccines, contain various components that might cause allergic reactions, including anaphylaxis. Not all such reactions are related to residual egg protein; however, the possibility of a reaction to influenza vaccines in egg-allergic people might be of concern to both these people and vaccine providers.

An egg-free recombinant influenza vaccine (RIV4, Flublok; Sanofi Pasteur) is available for people age 18 years and older and an egg-free cell culture-based IIV (cclIV4, Flucelvax; Seqirus) is approved for people age 4 years and older. ACIP does not state a preference for the use of RIV4 or cclIV4 for people with egg allergy although some providers may choose to administer RIV4 or cclIV4 to their patients with a history of severe egg allergy.

Reviews of studies of egg-culture based IIV and LAIV indicate that severe allergic reactions to egg-based influenza vaccines in people with egg allergy are unlikely. ACIP recommends that people with a history of egg allergy who have experienced only hives after exposure to egg may receive any recommended influenza vaccine (IIV, RIV4, LAIV4) appropriate for their age and health status.

In people with a history of severe egg allergy who report symptoms other than hives (e.g. angioedema, respiratory distress, recurrent vomiting) or who required emergent medical intervention (e.g., epinephrine) may also receive any recommended influenza vaccine appropriate for their age and health status. If a vaccine other than cclIV4 (Flucelvax) or RIV4 (Flublok) is used, it should be administered in a medical setting (e.g., a health department or physician office) and supervised by a healthcare provider who is able to recognize and manage severe allergic conditions. Providers should consider observing all patients for 15 minutes after vaccination to decrease the risk for injury should they experience syncope.

Inactivated influenza vaccines provided in multidose vials contain thimerosal as a preservative. Most people who had sensitivity to thimerosal when it was used in contact lens solution do not have reactions to thimerosal when it is used in vaccines. Check the package insert at [www.immunize.org/fda](http://www.immunize.org/fda) for a list of the vaccine components (i.e., excipients and culture media) used in the production of the vaccine, or go to [www.fda.gov/vaccines-blood-biologics/vaccines/vaccineslicensed-use-united-states](http://www.fda.gov/vaccines-blood-biologics/vaccines/vaccineslicensed-use-united-states).

For the 2020–2021 influenza season, no vaccine or packaging contains latex.

### 4. Has the person to be vaccinated ever had Guillain-Barré syndrome?

People who are not at high risk for severe influenza complications and who are known to have developed Guillain-Barré syndrome (GBS) within 6 weeks after receiving a previous influenza vaccination should not be vaccinated. As an alternative, clinicians might consider using influenza antiviral chemoprophylaxis for these people. Although data are limited, the established benefits of influenza vaccination for the majority of people who have a history of GBS, and who are at high risk for severe complications from influenza, justify yearly vaccination.